

Synthetic Communications

An International Journal for Rapid Communication of Synthetic Organic Chemistry

ISSN: 0039-7911 (Print) 1532-2432 (Online) Journal homepage: <http://www.tandfonline.com/loi/lcyc20>

A CONVENIENT METHOD FOR THE SYNTHESIS OF β -AMINO ACIDS VIA THE ARNDT-EISTERT APPROACH USING p-TOLUENESULPHONYL CHLORIDE AS A CARBOXYLIC GROUP ACTIVATING AGENT

Ganga-Ramu Vasanthakumar & Vommina V. Suresh Babu

To cite this article: Ganga-Ramu Vasanthakumar & Vommina V. Suresh Babu (2002) A CONVENIENT METHOD FOR THE SYNTHESIS OF β -AMINO ACIDS VIA THE ARNDT-EISTERT APPROACH USING p-TOLUENESULPHONYL CHLORIDE AS A CARBOXYLIC GROUP ACTIVATING AGENT, Synthetic Communications, 32:4, 651-657, DOI: [10.1081/SCC-120002414](https://doi.org/10.1081/SCC-120002414)

To link to this article: <http://dx.doi.org/10.1081/SCC-120002414>



Published online: 16 Aug 2006.



Submit your article to this journal [↗](#)



Article views: 86



View related articles [↗](#)



Citing articles: 5 View citing articles [↗](#)

Full Terms & Conditions of access and use can be found at
<http://www.tandfonline.com/action/journalInformation?journalCode=lcyc20>

**A CONVENIENT METHOD FOR
THE SYNTHESIS OF β -AMINO ACIDS
VIA THE ARNDT-EISTERT APPROACH
USING *p*-TOLUENESULPHONYL
CHLORIDE AS A CARBOXYLIC
GROUP ACTIVATING AGENT**

**Ganga-Ramu Vasanthakumar and
Vommina V. Suresh Babu***

Department of Studies in Chemistry,
Central College Campus, Bangalore University,
Bangalore 560 001, India

ABSTRACT


A simple method for the synthesis of Z-/Boc-/Fmoc-protected β -amino acids by the Arndt-Eistert approach employing *p*-toluenesulphonyl chloride for the activation of the carboxyl group of N^α -protected amino acid is described. The method is rapid and gave good yields with optical purity.

β -Amino acids exhibit a wide spectrum of biological importance similar to natural α -amino acids. They serve as building blocks for the synthesis of β -peptides,¹ and β -lactam antibiotics.² Seebach³ and Gellman^{4,5} groups reported that β -peptides also form stable secondary structures like their

*Corresponding author.

The use of *p*-toluenesulphonyl chloride (TsCl) in organic synthesis is widely known. It has been used for the activation of carboxylic group in the synthesis of amides and esters of simple carboxylic acids,¹¹ for the synthesis of peptides,¹² β -lactams¹³ etc. This paper describes the use of TsCl for the activation of carboxylic group in the synthesis of *N* $^{\alpha}$ -protected aminoacyldiazomethanes.

As shown in Scheme 1, N^α -protected aminoacyldiazomethane was prepared by addition of TsCl and diazomethane (CH_2N_2) in dichloromethane (CH_2Cl_2) to N^α -protected amino acid in THF at 0°C in presence of pyridine. The reaction proceeds through the formation of carboxylic-sulphonic mixed anhydride which was identified by IR spectra ($1825\text{--}1835\text{ cm}^{-1}$). The reaction was complete in about 30 min. All the resulting diazomethane derivatives were isolated as crystalline solids in good yield. They have been analysed by using IR by the presence of characteristic band at around $2105\text{--}2108\text{ cm}^{-1}$ ($-\text{COCHN}_2$ group) and $^1\text{H NMR}$. They were



Scheme 1. Synthesis of *N*-protected- β -homoamino acids.

then converted to their corresponding β -amino acids by the Wolff rearrangement using silver benzoate/1,4-dioxane/water by refluxing the mixture for 6 h at 70°C. All the *N*-protected β -amino acids were obtained as crystalline solids and were well characterised. The comparison of the determined optical rotations of the D- and L-isomers of Fmoc- α -Phg and that of β -Phg revealed that this method is completely free from racemisation.

Thus the present procedure is an efficient method for the synthesis of *N* $^{\alpha}$ -protected aminoacyldiazomethane using TsCl for carboxylic group activation. The advantages of this method are (a) the mild reaction conditions (b) the use of cheaper reagent (c) the general applicability (d) the selectivity, and (e) the good yields.

EXPERIMENTAL

All the amino acids used, unless otherwise specified has L-configuration only. Melting points reported are uncorrected. Thin layer chromatography (TLC) was carried out using the solvent systems: (i) ethyl acetate:hexane (35:65), (ii) chloroform:methanol:acetic acid (45:2:1), (iii) chloroform:methanol (9:1) and the R_f values are designated as R_{fA} , R_{fB} and R_{fC} respectively. ^1H NMR spectra were recorded on a Bruker AMX-400 MHz spectrometer. IR spectra were recorded on a Nicolet model impact 400 D FT-IR spectrometer. Elemental analyses were carried out by Perkin-Elmer Analyser and the samples were dried for 24 h under vacuum before analysis. Diazomethane gas was generated by heating *N*-nitroso-*N*-methyl-*p*-toluene sulphonamide with alcoholic KOH.¹⁴

General Procedure for Preparation of *N* $^{\alpha}$ -Protected Aminoacyldiazomethane Derivatives: TsCl (0.210 g, 1.1 mmol) and pyridine (0.08 mL, 1 mmol) were added to a solution of *N* $^{\alpha}$ -protected amino acid (1 mmol) in THF (5 mL) and stirred at 0°C for 15 min. The reaction mixture was treated with a saturated solution of CH_2N_2 in dry CH_2Cl_2 (20 mL) and stirred for 1 h at 0°C. The progress of the reaction was monitored by TLC and IR. After completion of the reaction, the excess of CH_2N_2 was decomposed by the dropwise addition of acetic acid. The mixture was washed with NaHCO_3 (25 mL \times 3), 5% HCl (25 mL \times 3) and brine, dried over anhydrous Na_2SO_4 and evaporated under reduced pressure. The resulting residue was recrystallized using CH_2Cl_2 /hexane to yield the title compound as a crystalline solid.

***N* $^{\alpha}$ -Fmoc-phenylalanyldiazomethane (IIa):** Yield, 94%; m.p., 136°C; R_{fA} , 0.62; R_{fB} , 0.8; $[\alpha]_D^{25} + 16.5$ ($c=1$, CHCl_3); Anal. Calc. for $\text{C}_{25}\text{H}_{23}\text{N}_3\text{O}_3$: C, 72.60; H, 5.60; N, 10.16; Found: C, 72.35; H, 5.57; N,



9.78; IR (γ_{\max}): 2108 cm^{-1} (-COCHN₂); ¹H NMR (CDCl₃, δ): 3.1 (2H, d), 4.2 (2H, m), 4.5 (2H, d), 5.2 (1H, s), 5.4 (1H, br) and 7.2–7.7 (13H, m).

N^α-Fmoc-valyldiazomethane (IIb): Yield, 85%; m.p., 124–125°C; *R*_fA, 0.69; *R*_fB, 0.81; [α]_D²⁵ – 23.40 (*c* = 1, CHCl₃); Anal. Calc. for C₂₁H₂₁N₃O₃: C, 69.4; H, 5.81; N, 11.56; Found: C, 69.28; H, 5.92; N, 11.46; IR (γ_{\max}): 2105 cm^{-1} (-COCHN₂); ¹H NMR (CDCl₃, δ): 0.90 (6H, d), 1.75 (1H, m), 4.10 (1H, m), 4.25 (1H, m), 4.45 (2H, m), 5.30 (1H, s), 5.4 (1H, d) and 7.3–7.9 (8H, m).

N^α-Fmoc-phenylglycinyldiazomethane (IIc): Yield, 90%; m.p., 148–149°C; *R*_fA, 0.66; *R*_fB, 0.80; [α]_D²⁵ – 32.3 (*c* = 1, CHCl₃); Anal. Calc. for C₂₄H₂₉N₃O₃: C, 72.50; H, 4.81; N, 10.57; Found: C, 72.68; H, 4.87; N, 10.6; IR (γ_{\max}): 2107 cm^{-1} (-COCHN₂); ¹H NMR (CDCl₃, δ): 4.25 (2H, m), 4.4 (2H, d), 5.1 (1H, s), 6.05 (1H, br) and 7.2–7.8 (13H, m).

N^α-Fmoc-D-phenylglycinyldiazomethane (IId): Yield, 92%; m.p., 153°C; *R*_fA, 0.7; *R*_fB, 0.83; [α]_D²⁵ + 32.3 (*c* = 1, CHCl₃); Anal. Calc. for C₂₄H₂₉N₃O₃: C, 72.50; H, 4.81; N, 10.57; Found: C, 72.43; H, 4.90; N, 10.52; IR (γ_{\max}): 2106 cm^{-1} (-COCHN₂); ¹H NMR (CDCl₃, δ): 4.2 (2H, m), 4.5 (2H, d), 5.1 (1H, s), 6.05 (1H, br) and 7.2–7.8 (13H, m).

N^α-Boc-alanyldiazomethane (IIe): Yield, 93%; m.p., 82°C; *R*_fA, 0.72; *R*_fB, 0.82; [α]_D²⁵ – 21.7 (*c* = 1, CHCl₃); Anal. Calc. for C, 50.69; H, 7.08; N, 19.70; Found: C, 50.80; H, 7.20; N, 19.47; IR (γ_{\max}): 2106 cm^{-1} (-COCHN₂); ¹H NMR (CDCl₃, δ): 1.05 (3H, d), 1.32 (9H, s), 4.21 (1H, m), 5.25 (1H, s) and 5.5 (1H, br).

N^α-Boc-valyldiazomethane (IIf): Yield, 90%; m.p., 64°C; *R*_fA, 0.72; *R*_fB, 0.83; [α]_D²⁵ – 30.4 (*c* = 1, CHCl₃); Anal. Calc. for C₁₁H₁₉N₃O₃: C, 54.75; H, 7.93; N, 17.41; Found: C, 54.62; H, 7.78; N, 17.42; IR (γ_{\max}): 2107 cm^{-1} (-COCHN₂); ¹H NMR (CDCl₃, δ): 0.92 (6H, d), 1.32 (9H, s), 1.75 (1H, m), 4.21 (1H, br), 5.20 (1H, s) and 5.42 (1H, br).

N^α-Z-phenylalanyldiazomethane (IIg): Yield, 92%; m.p., 82–83°C; *R*_fA, 0.68; *R*_fB, 0.78; [α]_D²⁵ + 48.2 (*c* = 1, CHCl₃); Anal. Calc. for C₁₈H₁₇N₃O₃: C, 66.86; H, 5.29; N, 12.99; Found: C, 66.72; H, 5.35; N, 12.86; IR (γ_{\max}): 2107 cm^{-1} (-COCHN₂); ¹H NMR (CDCl₃, δ): 2.9 (2H, d), 4.45 (1H, m), 5.0 (2H, s), 5.22 (1H, s), 5.46 (1H, br) and 7.2–7.3 (10H, m).

N^α-Z-alanyldiazomethane (IIh): Yield, 88%; m.p., 93–94°C; *R*_fA, 0.70; *R*_fB, 0.81; [α]_D²⁵ – 50.3 (*c* = 1, CHCl₃); Anal. Calc. for C₁₂H₁₃N₃O₃: C, 58.29; H, 5.29; N, 16.99; Found: C, 57.95; H, 5.13; N, 16.78; IR (γ_{\max}): 2107 cm^{-1} (-COCHN₂); ¹H NMR (CDCl₃, δ): 1.2 (3H, d), 4.26 (1H, m), 5.1 (2H, s), 5.25 (1H, s), 5.5 (1H, br) and 7.32 (5H, s).

N^α-Z-phenylglycinyldiazomethane (IIi): Yield, 86%; m.p., 83–84°C; *R*_fA, 0.69; *R*_fB, 0.79; [α]_D²⁵ – 36.4 (*c* = 1, CHCl₃); Anal. Calc. for C₁₇H₁₅N₃O₃: C, 66.01; H, 4.88; N, 13.58; Found: C, 65.72; H, 4.76; N, 13.72; IR



(γ_{\max}): 2108 cm^{-1} ($-\text{COCHN}_2$); $^1\text{H NMR}$ (CDCl_3 , δ): 4.5 (1H, br), 5.02 (2H, s), 5.22 (1H, s), 5.40 (1H, br) and 7.25 (10H, s).

General Procedure for Preparation of *N*-Protected β -Amino Acid: Silver benzoate (5.7 mg, 2.5×10^{-2} mmol) was added to a solution of *N* $^\alpha$ -protected aminoacyldiazomethane (1 mmol) in 1,4-dioxane (10 mL) and water (5 mL) and refluxed at 70°C for 6 h. It was filtered and the solvent was evaporated under reduced pressure. The residue was dissolved in saturated aqueous Na_2CO_3 (20 mL) and stirred for 20 min. The mixture was washed with ether (20 mL \times 3). The aqueous layer was acidified to pH 2 and extracted with ethyl acetate (20 mL \times 3). The extracts were pooled, washed with water (20 mL \times 2), dried over anhydrous Na_2SO_4 and evaporated under reduced pressure. The residue was precipitated using CH_2Cl_2 /hexane to get the title compound as a crystalline solid.

***N*-Fmoc- β -homophenylalanine (IIIa):** Yield, 85%; m.p., 110–112°C; $R_f\text{B}$, 0.63; $R_f\text{C}$, 0.75; $[\alpha]_D^{25} - 26.2$ ($c = 1$, CHCl_3); Anal. Calc. for $\text{C}_{25}\text{H}_{23}\text{NO}_4$: C, 75.05; H, 5.58; N, 3.70; Found: C, 75.21; H, 5.46; N, 3.80; IR (γ_{\max}): 1703 cm^{-1} (CO of urethane); $^1\text{H NMR}$ (CDCl_3 , δ): 2.43 (1H, d), 2.52 (2H, m), 2.71 (1H, d), 3.6 (1H, m), 4.1 (1H, m), 4.2 (2H, m) and 7.3–7.8 (14H, m).

***N*-Fmoc- β -homovaline (IIIb):** Yield, 80%; m.p., 153–154°C; $R_f\text{B}$, 0.60; $R_f\text{C}$, 0.78; $[\alpha]_D^{25} - 36.4$ ($c = 1$, CHCl_3); Anal. Calc. for $\text{C}_{21}\text{H}_{23}\text{NO}_4$: C, 70.14; H, 5.89; N, 4.20; Found: C, 70.06; H, 5.67; N, 4.35; IR (γ_{\max}): 1705 cm^{-1} (CO of urethane); $^1\text{H NMR}$ (CDCl_3 , δ): 0.85 (6H, d), 1.75 (1H, m), 2.30 (1H, d), 2.45 (1H, d), 3.75 (1H, m), 4.30 (3H, m) and 7.3–7.8 (9H, m).

***N*-Fmoc- β -homophenylglycine (IIIc):** Yield, 82%; m.p., 96–98°C; $R_f\text{B}$, 0.61; $R_f\text{C}$, 0.78; $[\alpha]_D^{25} - 22.2$ ($c = 1$, CHCl_3); Anal. Calc. for $\text{C}_{24}\text{H}_{21}\text{NO}_4$: C, 74.40; H, 5.45; N, 3.61; Found: C, 74.26; H, 5.23; N, 3.80; IR (γ_{\max}): 1705 cm^{-1} (CO of urethane); $^1\text{H NMR}$ (CDCl_3 , δ): 2.5 (2H, d), 4.25 (2H, m), 4.4 (2H, d), 5.85 (1H, br) and 7.2–7.8 (13H, m).

***N*-Fmoc- β -*D*-homophenylglycine (IIId):** Yield, 84%; m.p., 102°C; $R_f\text{B}$, 0.62; $R_f\text{C}$, 0.79; $[\alpha]_D^{25} + 21.8$ ($c = 1$, CHCl_3); Anal. Calc. for $\text{C}_{24}\text{H}_{21}\text{NO}_4$: C, 74.40; H, 5.45; N, 3.61; Found: C, 74.34; H, 5.16; N, 3.76; IR (γ_{\max}): 1705 cm^{-1} (CO of urethane); $^1\text{H NMR}$ (CDCl_3 , δ): 2.4 (2H, d), 4.25 (2H, m), 4.3 (2H, d), 5.80 (1H, br), 7.2–7.8 (13H, m) and 8.3 (1H, br).

***N*-Boc- β -homovaline (IIIe):** Yield, 88%; m.p., 98–99°C; $R_f\text{B}$, 0.62; $R_f\text{C}$, 0.72; $[\alpha]_D^{25} - 18.1$ ($c = 1$, CHCl_3); Anal. Calc. for $\text{C}_9\text{H}_{17}\text{NO}_4$: C, 53.45; H, 8.47; N, 6.92; Found: C, 53.28; H, 8.56; N, 6.56; IR (γ_{\max}): 1690 cm^{-1} (CO of urethane); $^1\text{H NMR}$ (CDCl_3 , δ): 1.2 (3H, d), 1.33 (9H, s), 2.4 (2H, d), 4.01 (1H, m), 5.25 (1H, br) and 8.4 (1H, br).

***N*-Boc- β -homovaline (IIIf):** Yield, 86%; m.p., 65–66°C; $R_f\text{B}$, 0.61; $R_f\text{C}$, 0.70; $[\alpha]_D^{25} - 20.2$ ($c = 1$, CHCl_3); Anal. Calc. for $\text{C}_{11}\text{H}_{21}\text{NO}_4$: C, 57.37; H, 9.19; N, 6.08; Found: C, 56.90; H, 9.34; N, 5.82; IR (γ_{\max}): 1690 cm^{-1}



(CO of urethane); ^1H NMR (CDCl_3 , δ): 0.94 (6H, d), 1.32 (9H, s), 1.72 (1H, m), 2.5 (2H, d), 4.20 (1H, br), 5.4 (1H, br) and 8.3 (1H, br).

N-Z- β -homophenylalanine (IIIg): Yield, 88%; m.p., 84–85°C; $R_f\text{B}$, 0.60; $R_f\text{C}$, 0.68; $[\alpha]_D^{25} - 36.2$ ($c = 1$, CHCl_3); Anal. Calc. for $\text{C}_{18}\text{H}_{19}\text{NO}_4$: C, 69.21; H, 6.13; N, 4.48; Found: C, 69.63; H, 6.47; N, 4.32; IR (γ_{max}): 1698 cm^{-1} (CO of urethane); ^1H NMR (CDCl_3 , δ): 2.36 (2H, d), 2.71 (2H, d), 3.8 (1H, m), 5.01 (2H, s), 5.3 (1H, m), 7.07 (5H, s), 7.28 (5H, s) and 9.4 (1H, br).

N-Z- β -homoolanine (IIIh): Yield, 84%; m.p., 122°C; $R_f\text{B}$, 0.60; $R_f\text{C}$, 0.69; $[\alpha]_D^{25} - 19.8$ ($c = 1$, CHCl_3); Anal. Calc. for $\text{C}_{12}\text{H}_{15}\text{NO}_4$: C, 61.00; H, 6.39; N, 5.92; Found: C, 61.27; H, 6.23; N, 5.87; IR (γ_{max}): 1700 cm^{-1} (CO of urethane); ^1H NMR (CDCl_3 , δ): 1.2 (3H, d), 2.4 (2H, d), 3.6 (1H, m), 5.1 (2H, s), 5.4 (1H, br), 7.31 (5H, s) and 8.4 (1H, br).

N-Z- β -homophenylglycine (IIIi): Yield, 82%; m.p., 86–88°C; $R_f\text{B}$, 0.63; $R_f\text{C}$, 0.74; $[\alpha]_D^{25} - 18.5$ ($c = 1$, CHCl_3); Anal. Calc. for $\text{C}_{17}\text{H}_{17}\text{NO}_4$: C, 68.44; H, 5.74; N, 4.95; Found: C, 68.23; H, 5.68; N, 4.67; IR (max): 1702 cm^{-1} (CO of urethane); ^1H NMR (CDCl_3 , δ): 2.5 (2H, d), 4.2 (1H, m), 5.01 (2H, s), 5.46 (1H, br), 7.25 (10H, s) and 8.6 (1H, br).

ACKNOWLEDGMENTS

We are grateful to Prof. K. M. Sivanandaiah and B. S. Sheshadri for usefull discussions, to Department of Science and Technology, Government of India for financial support. GRVK thanks KSVN Trust for their kind help.

REFERENCES

1. Banerjee, A.; Balaram, P. *Current Science* **1997**, *12*, 1067.
2. Kim, S.; Lee, P.H.; Lee, T.A. *J. Chem. Soc. Chem. Comm.* **1988**, 1242.
3. Seebach, D.; Matthews, J.L. *Chem. Commun.* **1997**, 2015.
4. Apella, D.H.; Christianson, L.A.; Klein, D.A.; Powell, D.R.; Huang, X.; Barchi Jr., J.J.; Gellman, S.H. *Nature* **1997**, *387*, 381.
5. Gellman, S.H. *Acc. Chem. Res.* **1998**, *31*, 173.
6. Hintermann, T.; Seebach, D. *Chimia* **1997**, *50*, 244.
7. Ellmerer-Muller, E.P.; Brossner, D.; Naslonh, N.; Tako, A. *Helv. Chim. Acta* **1998**, *81*, 59.
8. Leggio, A.; Liguori, A.; Procopio, A.; Sindona, G. *J. Chem. Soc. Perkin Trans. 1* **1997**, 1969.



SYNTHESIS OF β -AMINO ACIDS

657

9. Ananda, K.; Gopi, H.N.; Suresh Babu, V.V. *J. Peptide Research* **2000**, 55, 289.
10. (a) Suresh Babu, V.V.; Gopi, H.N.; Ananda, K. *J. Peptide Research* **1999**, 53, 308; (b) Ananda, K.; Suresh Babu, V.V. *Indian J. Chemistry* **1999**, 38B, 418.
11. Brewster, J.H.; Ciotti Jr., C.J. *J. Am. Chem. Soc.* **1955**, 77, 6214.
12. Theodoropoulos, D.; Gazopoulos, J. *J. Org. Chem.* **1962**, 27, 2091.
13. Wasserman, H.H.; Glazer, E.A.; Hearn, M.J. *Tetrahedron Lett.* **1973**, 49, 4855.
14. Furniss, B.S.; Hannaford, A.J.; Smith, P.W.G.; Tatchell, A.R. *Vogel's Textbook of Practical Organic Chemistry*, Educational Low-Priced Books Scheme, ELBS, Longman UK. Fifth Edition, **1989**, pp. 430–433.

Received in the Netherlands April 4, 2001



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

[Order now!](#)

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081SCC120002414>